

# **OCO (Orbiting Carbon Observatory) -2**

## **Spectroscopic needs for OCO-2**

---

**Revision Date: Dec 3, 2013**

**Jet Propulsion Laboratory  
Pasadena, California**

**Prepared By:**

Vivienne Payne, JPL  
OCO-2 ABSCO Lead

David Thompson, JPL  
Previous OCO-2 ABSCO Lead

### Revision History

Revision Date	Changes	Author
3 December 2013	Initial Release	V. Payne

## Table of Contents

<b>1</b>	<b>INTRODUCTION .....</b>	<b>5</b>
1.1	SCOPE AND BACKGROUND.....	5
1.2	DOCUMENT OVERVIEW.....	5
1.3	DATA USAGE POLICY.....	5
<b>2</b>	<b>ABSORPTION CROSS SECTION COMPUTATION .....</b>	<b>5</b>
<b>3</b>	<b>HDF5 FORMAT SPECIFICATION.....</b>	<b>9</b>
<b>4</b>	<b>REMAINING CHALLENGES AND FUTURE DIRECTIONS.....</b>	<b>10</b>
<b>5</b>	<b>REFERENCES .....</b>	<b>12</b>

# 1 Introduction

## 1.1 Scope and Background

This document is intended to provide an overview of the v4.2 ACOS/OCO-2 absorption coefficient (ABSCO) tables, key features and issues and potential directions for future research.

The ACOS/OCO-2 ABSCO tables are “lookup tables” that supply cross section values for absorbing gases in the Level 2 algorithm (L2) retrieval process. They record molecular absorption cross-sections over the range of relevant wavelengths, temperatures, and pressures in units of  $\text{cm}^2/\text{mol}$  for the gases  $\text{O}_2$ ,  $\text{H}_2\text{O}$ , and  $\text{CO}_2$ . During the L2 generation process the retrieval algorithm computes atmospheric absorption at each relevant temperature, pressure, and wavelength using linear interpolation. Successive versions have refined these tables by incorporating new laboratory results and theoretical models for increasingly accurate absorption coefficients. ABSCO tables are released together with each major build of the OCO-2 Level 2 algorithm. ABSCO v4.2 tables were used for the ACOS B3.5 Level 2 retrievals.

The tables can be computed at a spectral resolution of  $0.002 \text{ cm}^{-1}$  (required for validation of the tables using high resolution laboratory spectra) but the current L2 retrieval code uses a spectral resolution of  $0.01 \text{ cm}^{-1}$ .

## 1.2 Document Overview

Section 2 describes the absorption cross-section computation, while Section 3 describes the format of the ABSCO HDF files. Section 4 provides an overview of the remaining challenges and plans to address them.

## 1.3 Data Usage Policy


The OCO-2 ABSCO tables are provided freely to the public, on request. We request that when publishing using the tables, please acknowledge NASA and the ACOS/OCO-2 project.

- Include OCO-2 as a keyword to facilitate subsequent searches of bibliographic databases if it is a significant part of the publication
- Include a bibliographic citation for the tables. The most relevant citation currently is Thompson et al. (2012).
- We recommend sending courtesy copies of publications to the OCO-2 Project Scientist, Michael.R.Gunson@jpl.nasa.gov.

# 2 Absorption cross section computation

This section describes the studies providing line parameters for each of the main absorbing gases. Due to the modular nature of the ABSCO calculation, different bands rely on different codebases to compute the absorption cross sections. In summary, the LABFIT codebase from (Benner 2011) provides  $\text{CO}_2$  cross sections. This implements databases from (Benner 2011) and (Devi 2007) for  $2.0\mu\text{m}$  and  $1.6\mu\text{m}$   $\text{CO}_2$ , respectively. The procedure and modeling assumptions of this

approach are detailed at greater length in (Thompson 2012). A line mixing codebase from (Tran 2008) computes the major isotope of O<sub>2</sub>. We use the ABSCOI codebase, a routine adapted from the GFIT software package in use by the Total Column Carbon Observing Network (TCCON, Wunch 2010), for the minor isotopes of O<sub>2</sub> and all CH<sub>4</sub> and H<sub>2</sub>O lines. A flowchart showing the principal line list sources and the computation strategies for each line appears below in Figure 1.



**Figure 1. Absorption coefficient calculation processing and sources.**

In v4.2, the table format is a four-dimensional array to accommodate an additional “foreign broadener gas” dimension. This allows cross sections to be parameterized by temperature, pressure, and the volume mixing ratio of one other atmospheric gas. All tables are provided in this 4D format, though the calculations only incorporate broadening by H<sub>2</sub>O of O<sub>2</sub> and CO<sub>2</sub>. CH<sub>4</sub> tables using the HITRAN line list are now also available; they are computed over both CO<sub>2</sub> spectral windows. The temperature range has 17 levels and is intended to cover extremely low (terrestrial) temperatures. Table 1 below gives more detailed information on several key parameter sources.

Note that scaling factors are applied to the v4.2 absorption cross sections within the B3.5 Level 2 algorithm. O<sub>2</sub> cross sections are rescaled by a factor of 1.0125, CO<sub>2</sub> cross sections in the strong (2.06 μm) band are rescaled by 0.9946 and CO<sub>2</sub> cross-sections in the weak (1.6 μm) band are rescaled by 1.0038. The O<sub>2</sub> scaling factor was chosen in order to bring the mean retrieved surface pressure into agreement with ECMWF values. The CO<sub>2</sub> scaling factors were chosen in order to minimize disagreement with XCO<sub>2</sub> values from the Total Column Carbon Observing Network (TCCON) and to provide consistency between single-band CO<sub>2</sub> retrievals. The ABSCO HDF files themselves contain *unscaled* cross sections.

**Table 1: ABSCO v4.2 Parameter Sources**

	0.76 $\mu\text{m}$ O <sub>2</sub>	1.61 $\mu\text{m}$ CO <sub>2</sub>	2.06 $\mu\text{m}$ CO <sub>2</sub>	H <sub>2</sub> O
Spectral range	12745-13245 cm <sup>-1</sup>	4700-6500cm <sup>-1</sup>	4700-6500cm <sup>-1</sup>	12745-13245cm <sup>-1</sup> 4700-6500cm <sup>-1</sup>
Spectral resolution	0.01cm <sup>-1</sup> or 0.002cm <sup>-1</sup>	0.01 cm <sup>-1</sup> or 0.002cm <sup>-1</sup>	0.01 cm <sup>-1</sup> or 0.002cm <sup>-1</sup>	0.01 cm <sup>-1</sup> or 0.002cm <sup>-1</sup>
Position	Long (2010), Long (2011)	Devi (2007)	Benner/Devi (2011)	Gordon (2012), Rothman (2010)
Intensities	"	"	"	"
Air-widths	Tran (2008)	Predoi-Cross (2009)	"	"
Air-shifts	Brown (2009) Robichaud (2008a) Predoi-Cross (2008)	Devi (2007b)	"	"
Temp. dep.	Brown (2000)	Predoi-Cross (2009)	"	"
Line shapes	Voigt / Galatry	Speed-dependent Voigt	Speed-dependent Voigt	Voigt
Isotopologue abundance	Rothman (2009) <sup>1</sup>	Rothman (2009)	Rothman (2009)*	Rothman (2009)
H <sub>2</sub> O broadening	Drouin (2013)	Sung (2009)	Sung (2009)	-
Air-Line mixing	Tran (2008)	Devi (2007)	Benner/Devi (2011)	-
" Temp. dep.	Tran (2008)	-	-	-
Speed dep.	-	Devi (2007) <sup>1</sup>	Benner/Devi (2011)	-
Continuum	CIA via Tran (2008)	-	Mlawer (2012) *	Mlawer (2012) *

\*see text for further details

### 0.76 $\mu\text{m}$ O<sub>2</sub> Detail

The O<sub>2</sub> main isotopologue uses intensity and position parameters from new Cavity Ring Down Spectroscopy measurements (Long et al. 2010). The use of these parameters significantly reduces surface pressure error, while increasing yield.

Note that the (Long 2010) air-broadened half widths were *not* incorporated. The reason is that we found these parameters to be highly coupled with our line mixing strategy so that any change from the values provided in (Tran 2008) degraded performance. For consistency, we retain a Voigt line shape with widths and line mixing model exactly the same as in (Tran 2008). Those widths are based on a parameterization first found in (Yang 2005). Pressure shifts are from previous work by from (Robichaud 2008b) and (Predoi-Cross 2008) found in HITRAN 2008, with modifications to the R Branch by (Brown 2009). Minor transitions and isotopologues are computed using measurements from (Long 2011). The Galatry line shape was used for these minor lines, since narrowing parameters are available for lines in the Long et al. list. We

supplemented them with a few additional Voigt-profile lines from the HITRAN 2008 database to cover all remaining weak O<sub>2</sub> lines in the entire 12850-12900cm<sup>-1</sup> window.

Parameters for broadening of O<sub>2</sub> by H<sub>2</sub>O come from the study by Drouin et al. (2013).

Previous work (Butz 2010, Thompson 2012) has used an *ad hoc* rescaling of cross sections to reduce systemic surface pressure error. As stated previously, a rescaling factor of 1.0125 was used in this case to counter a surface pressure bias evidenced in the retrievals.

### 1.6μm and 2μm CO<sub>2</sub> Detail

The CO<sub>2</sub> bands use line parameters and mixing models derived from studies by Benner/Devi (2011) and Devi (2007). Additional detail on validation experiments for these databases, which are used in all line list versions after v4.0, are found in Thompson et al. (2012). The computation incorporates a speed dependent line profile with tridiagonal line mixing. Parameters for foreign broadening of CO<sub>2</sub> by H<sub>2</sub>O come from studies by Sung et al (2009). This study treated the 4.3μm band only. However, it is likely that these can be safely extrapolated to the other bands as well, and we favor this approach until additional direct laboratory data is available.

We modify the HITRAN 2008 isotopologue abundances slightly in the CO<sub>2</sub> band. These abundances are typical for measurements taken at sea level, but higher in the atmosphere CO<sub>2</sub> fractionation effects can modify the apparent abundances. Specifically, the 628 isotopologue abundance is increased by a factor of 1.04 to account for these apparent differences, and for the intrinsic uncertainty in band strength as measured in the laboratory. This factor was derived empirically from tests using the TCCON network of ground based, upward-looking FTS stations.

We note one final difference between the reference databases and our computed absorption coefficients. We found it necessary to incorporate an additional continuum absorption in the 2.06 μm CO<sub>2</sub> band which takes the shape of two Gaussian distributions centered at 4853.5 and 4789 wavenumbers, with respective standard deviations of 10 and 8 wavenumbers and intensity scalings of 2.1e-24 and 4.2e-25, respectively (this is the intensity at 1 Atm, but absorption at other levels scales in proportion to pressure). These parameters were set to minimize error in retrievals with TCCON uplooking FTS spectra.

### H<sub>2</sub>O Detail

We incorporate revised H<sub>2</sub>O line parameters from Gordon et al. [2012] for the 2μm region. It is likely that some or all of these parameters may become the standard for HITRAN 2012. We compute the H<sub>2</sub>O Continuum using the AER MT\_CKD codes (Mlawer et al., 2011). 1.67μm parameters are provided by HITRAN 2008, with the exception of a single line near 6290cm<sup>-1</sup> that we modified empirically by hand. This procedure, performed with the help of Dr. Iouli Gordon, reduced a prominent residual that we noted in both TCCON and GOSAT atmospheric retrievals. New line parameters from the HITEMP database (Rothman et al. 2010) have been used for weak water lines in the Oxygen A Band. Another change from the previous ABSCO version is that the line wing cutoff for H<sub>2</sub>O lines has been reduced to 25cm<sup>-1</sup> for consistency with the assumptions of the MT\_CKD continuum model. This was a minor error in previous versions, and it is unlikely to have significantly impacted retrieval results. Finally, the continuum itself has been revised. We use an unofficial modified version of the AER MT\_CKD continuum (Mlawer et al. 2012), supplied for testing purposes only by Eli Mlawer. This increases the continuum slightly



over the previous version of the AER MT\_CKD model in the OCO-2 spectral intervals. There is no absolute consensus yet on the continuum level in the 2.06 micron band. This revision offers a conservative compromise between the previous continuum and new measurements by Ptashnik et al. (2011). The revised MT\_CKD also falls relatively close to new measurements by Mondelain et al. (2013).

Overall we find that the combination of these changes improves the spectrum fit quality for typical TCCON and GOSAT retrievals.

### 3 HDF5 Format Specification

The ABSCO tables are provided in an HDF5 format. The table is a 4D structure indexed in the following order:

- pressure levels 1 to  $N_{\text{pres}}$
- temperature levels from 1 to  $N_{\text{temps}}$
- foreign broadener vmrs from 1 to  $N_{\text{vmrs}}$
- frequencies from 1 to  $N_{\text{freqs}}$

In general, each table models one absorbing gas and up to one foreign broadening gas (here, always H<sub>2</sub>O). We refer to the absorber's HITRAN gas index value with the symbol  $Q_{\text{abs}}$  and its HITRAN isotope index as  $R_{\text{abs}}$ . The foreign broadener index is  $Q_{\text{brd}}$ .

The HDF file is has the following format:

- **File:** The top-level file object represents a collection of one or more absorber gas cross sections that have been computed for a common spectral range and set of atmospheric conditions. The file contains top-level attributes:
  - version: a version string describing the table release
  - addl\_ident: a special identifier string, if any
  - gas\_name: a string such as "o2"
  - wn\_begin: the starting wavenumber
  - wn\_end: the ending wavenumber
  - comment: a space for notes about the table creation
- **Gas\_ $[Q_{\text{abs}}]$ \_Absorption** The absorption coefficients are a 4D table of size  $N_{\text{pres}} \times N_{\text{temps}} \times N_{\text{vmrs}} \times N_{\text{freqs}}$  and are indexed in that order. The dataset attributes are:
  - addl\_ident: a special identifier string, if any
  - gas\_name: a string such as "o2", which should match its filename
  - comment: a space for notes
- **Gas\_Index** a string containing the 2 digit HITRAN index of the principal absorbing gas, equivalent to  $Q_{\text{abs}}$  above.
- **Pressure** The pressure is a dataset of size  $N_{\text{pres}}$  representing the pressure in Pascals at each atmospheric level in the table
- **Temperature** Temperature is a 2D dataset of degrees Kelvin, of shape  $N_{\text{pres}} \times N_{\text{temps}}$ . It records the temperature grid point values, which might differ depending on pressure level.
- **Broadener\_ $[Q_{\text{brd}}]$ \_VMR** A dataset of size  $N_{\text{vmrs}}$  representing the different volume mixing ratios at which the foreign broadening gas is modeled.  $Q_{\text{brd}}$  is a HITRAN index. Now we only model H<sub>2</sub>O broadening, so the index is always 01.

- **broadener\_name**: the string “h2o”
- **Broadener\_Index** a string contraining the 2 digit HITRAN index of the broadening gas, equivalent to  $Q_{brd}$  above
- **Wavenumber** The Wavenumber object is a dataset of size  $N_{freqs}$  describing the frequency grid spacing.

## 4 Remaining challenges and future directions

The goal for the OCO-2 mission is to achieve 1 ppm accuracy in the XCO<sub>2</sub> retrievals. In order to allow room for other terms in the error budget, the aim is to achieve 0.1 % accuracy in the spectroscopic input to the forward model.

Figure 2, Figure 3 and Figure 4 show the magnitude of remaining residuals in the O<sub>2</sub> A band, the weak CO<sub>2</sub> band and the strong CO<sub>2</sub> band respectively. (Strictly speaking, the “current” residuals in these Figures are residuals using ABSCO v4.0. Residuals for ABSCO v4.2 are visually very similar, although the updates between v4.0 and v4.2 did result in small changes in the Level 2 retrieval results.) The differences between “previous” and “current” in these Figures are described in more detail in Thompson et al. (2012). All three bands show evidence of systematic spectral residuals with the current ABSCO version.






Figure 2. Averaged GOSAT residuals for the 0.76  $\mu\text{m}$  O<sub>2</sub> A-band.



**Figure 3. Averaged GOSAT residuals for the weak ( $1.6\ \mu\text{m}$ )  $\text{CO}_2$  band. (Figure from Thompson et al., 2012)**



**Figure 4. Averaged GOSAT residuals for the strong ( $2.06\ \mu\text{m}$ )  $\text{CO}_2$  band. (Figure from Thompson et al., 2012)**

It has been shown (e.g. Thompson et al., 2012) that the traditionally used Voigt lineshape does not provide the required accuracy in this region. Research into the “best” lineshape is an ongoing

endeavor within the spectroscopy community. The approach taken by the OCO-2 ABSCO team has been to work towards simultaneous fitting of multiple laboratory spectra (Benner and Devi, 2011; Devi et al., 2007), where the laboratory conditions cover the full range of conditions encountered in Earth's atmosphere, in order to provide a consistent set of spectroscopic parameters for OCO-2.

Work is underway to apply the multi-spectrum fitting approach to FTS, CRDS and Photoacoustic Spectroscopy (PAS) measurements in the O<sub>2</sub> A band. The multi-spectrum fitting approach is expected to offer significant improvement over our current ABSCO version and to provide new insight into lineshape and collision-induced absorption (CIA) in the A band.

To date, this multi-spectrum fitting approach has been applied to room temperature Fourier Transform Spectrometer (FTS) laboratory measurements of CO<sub>2</sub> spectra. Resulting improvements in fits to atmospheric measurements are described in Thompson et al. (2012). However, the updated fits still do not allow us to fit available measurements within the noise. Work is underway to incorporate new cold temperature FTS measurements from JPL as well as Cavity Ringdown Spectroscopy (CRDS) measurements from NIST and Caltech into these fits. The new fits are expected to place further constraints on the spectroscopic parameters and lineshape in the OCO-2 CO<sub>2</sub> bands.

It will be necessary to continue to assess the need for laboratory measurements that can fill the gaps in our existing knowledge, to incorporate progress in theoretical studies into the multi-spectrum fitting approach and to continue to validate the ABSCO tables against high quality laboratory and atmospheric measurements (with various viewing geometries).

## 5 References

- [Benner/Devi 2011] Benner DC, Devi VM, Nugent E, Sung K, Brown LR, Miller CE. et al., Line parameters of carbon dioxide in the 4850 cm<sup>-1</sup> region. In: Twenty-second Coll. on High resolution molecular spectroscopy, Dijon, Fr 2011; poster N19.
- [Brown 2000] Brown, L. R., Plymate, C. Experimental line parameters of the oxygen A band at 760nm. J Mol Spectrosc 2000; 199:166–79.
- [Brown 2009] Line list is available in the acos/absco repository at the following location: acos/level\_2/exe/absco/input/line\_mixing/o2/lm\_o2\_inputs\_20091026/SDF.dat
- [Devi 2007] Devi, V. M., D. C. Benner, L. R. Brown, C. E. Miller, and R. A. Toth (2007), Line mixing and speed dependence in CO<sub>2</sub> at 6227.9 cm<sup>-1</sup>: Constrained multispectrum analysis of intensities and line shapes in the 30013 ← 00001 band, J. Mol. Spectrosc. 245, 52-80.
- [Dudhia 2009] Anu Dudhia, RFM Software package and user guide available at <http://www.atm.ox.ac.uk/RFM/>.
- [Drouin 2013] B.J. Drouin, V. H. Payne, F. Oyafuso, K. Sung and E. J. Mlawer, Pressure broadening of oxygen by water, J. Quant. Spectrosc. & Radiat. Trans., in press, 2013.
- [Gordon 2012] I.E. Gordon, L. Lodi, G.C. Toon, J. Tennyson, L.R. Brown, L.S. Rothman. "Evaluation of the experimental and theoretical spectral parameters of water vapor lines in the 2-μm region using Solar-pointing FTS spectra," (2012) in preparation.
- [Hartmann 2009] J.-M. Hartmann, H. Tran, and G. C. Toon, *Influence of line mixing on the retrievals of atmospheric CO<sub>2</sub> from spectra in the 1.6 and 2.1 μm regions*, Atmos. Chem. Phys. Discuss., 9, 1–26, 2009
- [Lamoureux 2010] J. Lamoureux, H. Tran, A.L. Laraia, R.R. Gamache, L.S. Rothman, I.E. Gordon, and J.-M. Hartmann, J. Updated database plus software for line-mixing in CO<sub>2</sub> infrared spectra and their test using laboratory spectra in the 1.5-2.3 μm region, Quant. Spectrosc. and Rad. Transfer 111, 2321-2331 (2010).

- [Long 2010] D.A. Long, D.K. Havey, M. Okumura, C.E. Miller, J.T. Hodges, O<sub>2</sub> A-band line parameters to support atmospheric remote sensing, *Journal of Quantitative Spectroscopy and Radiative Transfer*, Volume 111, Issue 14, September 2010, Pages 2021-2036, ISSN 0022-4073, 10.1016/j.jqsrt.2010.05.011.
- [Mlawer 2012] Mlawer, E.J., V.H. Payne, J.-L. Moncet, J.S. Delamere, M.J. Alvarado, D.D. Tobin, Development and recent evaluation of the MT\_CKD model of continuum absorption, submitted to *Phil. Trans. Roy. Soc.A*, June 13, 2012 370 1968 2520-2556.
- [Mondelain 2013] D. Mondelain, A. Aradj, S. Kassi and A. Campargue, The water vapor self continuum by CRDS at room temperature in the 1.6 mm transparency window, JQSRT-D-13-00183R1, in press
- [Niro 2004] Niro, F., Boulet, C., and Hartmann, J. M.: Spectra calculations in central and wing regions of CO<sub>2</sub> IR bands between 10 and 20  $\mu$ m. I: Model and laboratory measurements, *J. Quant. Spectrosc. Radiat. Transfer* 88, 483-498, 2004.
- [Niro 2005] F. Niro, T. von Clarmann, K. Jucks, and J.-M. Hartmann (2005), Spectra calculations in central and wing regions of CO<sub>2</sub> IR bands between 10 and 20  $\mu$ m. III: atmospheric emission Spectra, *J. Quant. Spectrosc. Radiat. Transfer* 90, 61-76.
- [Predoi-Cross 2008] Predoi-Cross, A., C. Holladay, H. Heung, J.-P. Bouanich, G. Ch. Mellau, R. Keller, and D. R. Hurtmans (2008), Nitrogen-broadened lineshapes in the oxygen A-band: Experimental results and theoretical calculations, *J. Mol. Spectrosc.* 251, 159-175.
- [Predoi-Cross 2009] Predoi-Cross, A., A. R. W. McKellar, D. Chris Benner, V. Malathy Devi, R. R. Gamache, C. E. Miller, R. A. Toth, and L. R. Brown (2009), Temperature dependences for air-broadened Lorentz half width and pressure-shift coefficients in the 30013-00001 and 30012-00001 bands of CO<sub>2</sub> near 1600 nm, *Can. J. Phys.* (in press).
- [Ptashnik 2011] Ptashnik et al. *Journal of Geophysical Research*, vol. 116, D16305, doi:10.1029/2011JD015603, 2011.
- [Robichaud 2008a] Robichaud, D. J., Hodges, J. T., Maslowski, P., Yeung, L.Y., Okumura, M., Miller, C. E., et al. High-accuracy transition frequencies for the O<sub>2</sub> A-band. *J Mol Spectrosc* 2008;251:27-37.
- [Robichaud 2008b] Robichaud, D. J., Hodges, J. T., Brown, L. R., Lisak, D., Maslowski, P., Yeung, L. Y., et al. Experimental intensity and line shape parameters of the oxygen A-band using frequency-stabilized cavity ring-down spectroscopy. *J Mol Spectrosc.* 2008; 248:1-13.
- [Robichaud 2008c] Robichaud D. J., Hodges, J. T., Lisak, D., Miller, C. E., Okumura, M. High-precision pressure shifting measurement technique using frequency-stabilized cavity ring-down spectroscopy. *JQSRT* 2008;109:435-44.
- [Rothman 2005] Rothman, L. S., et al. (2005), The HITRAN molecular spectroscopic database: Edition of 2004, *J. Quant. Spectrosc. Radiat. Transfer*, 96, 139 - 204.
- [Rothman 2009] Rothman, L. S. et al. *The HITRAN 2008 molecular spectroscopic database*. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 110 (2009) 533-572.
- [Rothman 2010] Rothman, L. S., I. E. Gordon, R. J. Barber, H. Dothe, R. R. Gamache, A. Goldman, V. I. Perevalov, S. A. Tashkun, J. Tennyson (2010). HITRAN, the high-temperature molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer*. 111:15, p. 2139-2150.
- [Sung 2009] Keeyoon Sung; L. R. Brown, R. A. Toth, T. J. Crawford. Fourier transform infrared spectroscopy measurements of H<sub>2</sub>O-broadened half-widths of CO<sub>2</sub> at 4.3  $\mu$ m. *Canadian Journal of Physics*; May 2009, Vol. 87 Issue 5, p. 469.
- [Thompson 2012] Thompson, D. R., D. C. Benner, L. R. Brown, D. Crisp, V. Malathy Devi, Y. Jiang, V. Natraj, F. Oyafuso, K. Sung, D. Wunch, R. Castaño, C. E. Miller Atmospheric validation of high accuracy CO<sub>2</sub> absorption coefficients for the OCO-2 mission. *Journal of Quantitative Spectroscopy and Radiative Transfer* (2012), <http://dx.doi.org/10.1016/j.jqsrt.2012.05.021>.
- [Tran 2006] Tran, H, Boulet, C, and Hartmann, J.-M. Line mixing and collision-induced absorption by oxygen in the A band: Laboratory measurements, model, and tools for atmospheric spectra computations. *Journal of Geophysical Research*, vol. 111, D15210, doi:10.1029/2005JD006869, 2006.
- [Tran 2008] Tran, H. and Hartmann, J. M. An improved O<sub>2</sub> A band absorption model and its consequences for retrievals of photon paths and surface pressures, *Journal of Geophysical Research*, vol 113, D18104, 2008.
- [Toth 2007] Toth, R. A., C. E. Miller, V. M. Devi, D. C. Benner, and L. R. Brown (2007), Air-broadened width and pressure shift coefficients of 12C16O<sub>2</sub>: 4700 - 7000 cm<sup>-1</sup>, *J. Mol. Spectrosc.* 246, 133-157.
- [Toth 2008] Toth, R. A., Brown, L. R., Miller, C. E., Malathy Devi, V., and Benner, D. C.: Spectroscopic database of CO<sub>2</sub> line parameters: 4300-7000 cm<sup>-1</sup>, *J. Quant. Spectrosc. Radiat. Transfer* 109, 906-921, 2008.
- [Toon 2010] Toon, G. Personal Communication.

- [Vess 2012] E. M. Vess, C. J. Wallace, H. M. Campbell, V. E. Awadalla, J. T. Hodges, D. A. Long, and D. K. Havey. Measurement of H<sub>2</sub>O Broadening of O<sub>2</sub> A-Band Transitions and Implications for Atmospheric Remote Sensing. *Journal of Physical Chemistry*. 2012 (in press).
- [Wunch 2010], Wunch, D., Toon, G. C., Wennberge, P. O. et al. Calibration of the total carbon column observing network using aircraft profile data. *Atmos Meas. Tech. Discuss.*, 3, 2603-2632, 2010. [www.atmos-meas-tech-discuss.net/3/2603/2010/](http://www.atmos-meas-tech-discuss.net/3/2603/2010/) doi:10.5194/amtd-3-2603-2010